

Dynamical modeling of microRNA action on the protein translation process

Andrei Zinovyev^{1,2,3,6,*}, Nadya Morozova⁴, Nora Nonne⁴
Emmanuel Barillot^{1,2,3}, Annick Harel-Bellan⁴, Alexander N. Gorban^{5,6}

¹ Institut Curie, Service Bioinformatique, Paris, France

² INSERM, U900, Paris, F-75248 France

³ Ecole des Mines de Paris, ParisTech, Fontainebleau, France

⁴ CNRS FRE 2944, Institut André Lwoff, Villejuif, France

⁵ University of Leicester, UK

⁶ Institute of Computational Modeling SB RAS, Krasnoyarsk, Russia

* Correspondent author e-mail: andrei.zinovyev@curie.fr

November 9, 2009

Abstract

Protein translation is a multistep process which can be represented as a cascade of biochemical reactions (initiation, ribosome assembly, elongation, etc.), the rate of which can be regulated by small non-coding microRNAs through multiple mechanisms. It remains unclear what mechanisms of microRNA action are most dominant: moreover, many experimental reports deliver controversial messages on what is the concrete mechanism actually observed in the experiment. Parker and Nissan [37] demonstrated that it is impossible to distinguish alternative biological hypotheses using the steady state data on the rate of protein synthesis. For their analysis they used two simple kinetic models of protein translation. In contrary, we show that dynamical data allow to discriminate some of the mechanisms of microRNA action. We demonstrate this using the same models as in [37] for the sake of comparison but the methods developed (asymptotology of biochemical networks) can be used for other models. As one of the results of our analysis, we formulate a hypothesis that the effect of microRNA action is measurable and observable only if it affects the dominant system (generalization of the limiting step notion for complex networks) of the protein translation machinery. The dominant system can vary in different experimental conditions that can partially explain the existing controversy of some of the experimental data.

1 Introduction

MicroRNAs (miRNAs) are currently considered as key regulators of a wide variety of biological pathways, including development, differentiation and oncogenesis. Recently, remarkable progress was made in the understanding of microRNA biogenesis, function and mechanism of action. Mature microRNAs are incorporated into the RISC effector complex, which includes as a key component an Argonaute protein. MicroRNAs affect gene expression by guiding the RISC complex toward specific target mRNAs. The exact mechanism of this inhibition is still a matter of debate. In the past few years, several mechanisms have been reported, some of which are contradictory (for review, see [6, 9, 11]). These include in particular inhibition of translation initiation (acting at the level of cap-40S or 40S-AUG-60S association steps), inhibition of translation elongation or premature termination of translation. MicroRNA-mediated mRNA decay and sequestration of target mRNAs in P-bodies have been also proposed. Moreover, some microRNAs mediate target mRNA cleavage [56], chromatin reorganization followed by transcriptional repression or activation [26, 41], or translational activation [40, 50].

The most frequently reported, but also much debated, mechanism of gene repression by microRNAs occurs at the level of mRNA translation. This is particularly true when the microRNA is not fully complementary to its target. At this level, several mode of actions have been suggested (see Fig. 1). Historically, the first proposed mechanism was inhibition of translation elongation. The major argument supporting this hypothesis was the observation that the inhibited mRNA remained associated with the polysomal fraction (in which mRNAs are associated with polysomes), an observation that was reproduced in different systems [18, 38, 43, 39, 35]. The idea of a post-initiation mechanism was further supported by the observation that some mRNAs can be repressed by a microRNA even when their translation is cap-independent (mRNAs with an IRES or A-capped) [1, 25, 33, 43]. Although it was initially proposed that the ribosomes was somehow “frozen” on the mRNA, it is important to note that it is difficult to discriminate experimentally between different post-initiation potential mechanisms, such as elongation inhibition, premature ribosome dissociation (“ribosome drop-off”) or normal elongation but nascent polypeptide degradation. The last proposition (which can occur in conjunction with the two others) is supported by the fact that the mRNA-polysomal association is puromycin-sensitive, indicating that it depends on a peptidyl-transferase activity [2, 35]. However, no nascent peptide has ever been experimentally demonstrated; thus degradation would occur extremely rapidly after synthesis [42, 43, 38]. Degradation, in any case, is proteasome-independent [42]. Premature ribosome dissociation is supported by decreased read-through of inhibited mRNA [43]. Ribosome drop-off and/or ribosomal “slowing” are supported by the slight decrease in the number of associated ribosomes observed in some studies [38, 35].

Concurrently, several reports have been published indicating an action of microRNAs at the level of initiation. An increasing number of papers reports that microARN-targetted mRNAs shift toward light fractions in polysomal profiles [3, 27, 42]. This shows a decrease of mature translating ribosomes, suggesting that microRNAs act on the initiation step. Moreover, several reports show that microRNA-mediated inhibition is relieved when translation is driven by a cap-independent mechanism such as IRES-mRNA or A-capped-mRNA [20, 27, 42]. This observation was confirmed in several in-vitro studies [36, 48, 53, 51]. In particular, in one of those, an excess of eIF4F could relieve the inhibition, and inhibition lead to decreased 80S in polysomal gradient [36].

Most of the data indicating a shift toward light polysomal fraction or requirement for a cap-dependent translation are often interpreted in favour of involvement of microRNAs at early steps of translation, i.e., cap binding and 40S recruitment. However, some of them are also compatible with a block at the level of 60S subunit joining. This hypothesis is also supported by in-vitro experiments showing a lower amount of 60S relative to 40S on inhibited mRNAs. Moreover, toe-printing experiments show that 40S is positioned on the AUG [52]. Independently, it was shown that eIF6, an inhibitor of 60S joining, is required for microRNA action [7], but this was contradicted by other studies [9].

Thus, data regarding the exact step of translational inhibition are clearly contradictory. Taking also into account the data about mRNA degradation and P-bodies localization, it is difficult to draw a clear picture of the situation, and the exact mechanism by which microRNA repress mRNA expression is highly controversial, let alone the interrelations between the different mechanisms and their possible concomitant action. Several attempts to integrate the different hypothesis have been made [6, 9, 11, 21, 29, 49]. For example, one mechanism could act as a “primary” effect, and the other as a “secondary” mechanism, either used to reinforce the inhibition or as back-up mechanism. In others, the different mechanisms could all coexist, but occur differentially depending on some yet unidentified characteristic. For example, it has been observed than the same mRNA targeted by the same microRNA can be regulated either at the initiation or the elongation step depending on the mRNA promoter and thus on mRNA nuclear history [28]. It was also proposed that technical (experimental) problems, including the variety of experimental systems used, may also account for these discrepancies [6, 9, 11]. However, this possibility does not seem to be sufficient to provide a simple and convincing explanation to the reported discrepancies.

A possible solution to exploit the experimental observations and to provide a rational and straightforward interpretation is the use of mathematical models for microRNA action. Nissan and Parker recently studied the steady states of two simple kinetic models [37] and provided critical analysis for the experiments with alternative mRNA cap structures and IRES elements [36, 48, 51], leading to possible explanation of the conflicting results. The authors suggested that the relief of translational repression upon replacement of the cap structure can be explained if microRNA is acting on a step which is not rate-limiting in the modified system, in which case, an effect of microRNA can simply




Figure 1: Interaction of microRNA with protein translation process. Four mechanisms of translation repression which are considered in the mathematical modeling are indicated: 1) on the initiation process, preventing assembling of the initiation complex; 2) on a late initiation step, such as searching for the start codon; 3) on the ribosome assembly; 4) on the translation process. There exist other mechanisms of microRNA action on protein translation (transcriptional, transport to P-bodies, ribosome drop-off, co-translational protein degradation and others) that are not considered in this paper. Here 40S and 60S are light and heavy components of the ribosome, 80S is the assembled ribosome bound to mRNA, eIF4F is an translation initiation factor, PABPC1 is the Poly-A binding protein, “cap” is the mRNA cap structure needed for mRNA circularization, RISC is the RNA-induced silencing complex.

not be observed. It was claimed that it is impossible to discriminate between two alternative interpretations of the biological experiments with cap structure replacement from the sole monitoring of the steady state level of protein [37].

Two remarks can be made in this regard. First, in practice not only the steady state level of protein can be observed but also other dynamical characteristics, such as the *relaxation time*, i.e. the time needed to achieve the steady state rate after a perturbation (such as restarting the translation process). We argue that having these measurements in hands, *one can distinguish between two alternative interpretations*. In this paper we provide such a method from the same models as constructed by Nissan and Parker, for comparison purposes. However, the method applied can be easily generalized for other models.

Second, even in the simple non-linear model of protein translation, taking into account the recycling of ribosomal components, it is difficult to define what is the rate limiting step. It is known from the theory of asymptotology of biochemical networks [17] that even in complex linear systems the “rate limiting place” notion is not trivial and can not be reduced to a single reaction step. Moreover, in non-linear systems the “rate limiting place” can change with time and depend on the initial conditions. Hence, conclusions of [37] should be re-considered for the non-linear model, made more precise and general. The notion of rate limiting step should be replaced by the notion of *dominant system*.

In this paper we perform careful analysis of the Nissan and Parker’s models and provide their approximate analytical solutions, which allows us to generalize the conclusions of [37] and make new testable predictions on the identifiability of active mechanism of microRNA-dependent protein translation inhibition.

2 Methods

2.1 Asymptotology and dynamical limitation theory for biochemical reaction networks

Most of mathematical models that really work are simplifications of the basic theoretical models and use in the backgrounds an assumption that some terms are big, and some other terms are small enough to neglect or almost neglect them. The closer consideration shows that such a simple separation on “small” and “big” terms should be used with precautions, and special culture was developed. The name “asymptotology” for this direction of science was proposed by [30] defined as “the art of handling applied mathematical systems in limiting cases”.

In chemical kinetics three fundamental ideas were developed for model simplification: *quasiequilibrium asymptotic* (QE), *quasi steady-state asymptotic* (QSS) and the idea of *limiting step*.

In the IUPAC Compendium of Chemical Terminology (2007) one can find a definition of limiting step [44]: “A rate-controlling (rate-determining or rate-limiting) step in a reaction occurring by a composite reaction sequence is an elementary reaction the rate constant for which exerts a strong effect – stronger than that of any other rate constant – on the overall rate.”

Usually when people are talking about limiting step they expect significantly more: there exists a rate constant which exerts such a strong effect on the overall rate that the effect of all other rate constants together is significantly smaller. For the IUPAC Compendium definition a rate-controlling step always exists, because among the control functions generically exists the biggest one. On the contrary, for the notion of limiting step that is used in practice, there exists a difference between systems with limiting step and systems without limiting step.

During XX century, the concept of the limiting step was revised several times. First simple idea of a “narrow place” (the least conductive step) could be applied without adaptation only to a simple cycle or a chain of irreversible steps that are of the first order (see Chap. 16 of the book [23] or the paper by [5] or the section 2.5 of this paper). When researchers try to apply this idea in more general situations they meet various difficulties.

Recently, we proposed a new theory of dynamic and static limitation in multiscale reaction networks [16, 17]. This approach allows to find the simplest network which can substitute a multiscale reaction network such that the dynamics of the complex network can be approximated by the simpler one. Following the asymptotology terminology [54], we call this simple network the *dominant system* (DS). In the simplest cases, the dominant system is a subsystem of the original model. However, in the general case, it also includes new reactions with kinetic rates expressed through the parameters of the original model, and rates of some reactions are renormalized: hence, *in general, the dominant system is not a subsystem of the original model*.

The dominant systems can be used for direct computation of steady states and relaxation dynamics, especially when kinetic information is incomplete, for design of experiments and mining of experimental data, and could serve as a robust first approximation in perturbation theory or for preconditioning. Dominant systems serve as correct generalization of the limiting step notion in the case of complex multiscale networks. They can be used to answer an important question: given a network model, which are its critical parameters? Many of the parameters of the initial model are no longer present in the dominant system: these parameters are non-critical. Parameters of dominant systems indicate putative targets to change the behavior of the large network.

Most of reaction networks are nonlinear, it is nevertheless useful to have an efficient algorithm for solving linear problems. First, nonlinear systems often include linear subsystems, containing reactions that are (pseudo)monomolecular with respect to species internal to the subsystem (at most one internal species is reactant and at most one is product). Second, for binary reactions $A + B \rightarrow \dots$, if concentrations of species A and B (c_A, c_B) are well separated, say $c_A \gg c_B$ then we can consider this reaction as $B \rightarrow \dots$ with rate constant proportional to c_A which is practically constant, because its relative changes are small in comparison to relative changes of c_B . We can assume that this condition is satisfied for all but a small fraction of genuinely non-linear reactions (the set of non-linear reactions changes in time but remains small). Under such an assumption, non-linear behavior can be approximated as a sequence of such systems, followed one each other in a sequence of “phase transitions”. In these transitions, the order relation between some of species concentrations changes. Some applications of this approach to systems biology are presented by [45]. The idea of controllable linearization

“by excess” of some reagents is in the background of the efficient experimental technique of Temporal Analysis of Products (TAP), which allows to decipher detailed mechanisms of catalytic reactions [55].

Below we give some details on the approaches used in this paper to analyse the models by Nissan and Parker [37].

2.2 Notations

To define a chemical reaction network, we have to introduce:

- a list of components (species);
- a list of elementary reactions;
- a kinetic law of elementary reactions.

The list of components is just a list of symbols (labels) $A_1, \dots, A_n$. Each elementary reaction is represented by its *stoichiometric equation*

$$\sum_{si} \alpha_{si} A_i \rightarrow \sum_{si} \beta_{si} A_i, \quad (1)$$

where s enumerates the elementary reactions, and the non-negative integers α_{si} , β_{si} are the *stoichiometric coefficients*. A stoichiometric vector γ_s with coordinates $\gamma_{si} = \beta_{si} - \alpha_{si}$ is associated with each elementary reaction.

A non-negative real *extensive* variable $N_i \geq 0$, amount of A_i , is associated with each component A_i . It measures “the number of particles of that species” (in particles, or in moles). The concentration of A_i is an *intensive* variable: $c_i = N_i/V$, where V is volume. In this paper we consider the volume (of cytoplasm) to be constant. Then the kinetic equations have the following form

$$\frac{dc}{dt} = \sum_s w_s(c, T) \gamma_s + v, \quad (2)$$

where T is the temperature, w_s is the rate of the reaction s , v is the vector of external fluxes normalized to unite volume. It may be useful to represent external fluxes as elementary reactions by introduction of new component $\emptyset$ together with income and outgoing reactions $\emptyset \rightarrow A_i$ and $A_i \rightarrow \emptyset$.

The most popular *kinetic law* of elementary reactions is the *mass action law* for perfect systems:

$$w_s(c, T) = k_s \prod_i c_i^{\alpha_{si}}, \quad (3)$$

where k_s is a “kinetic constant” of the reaction s .

2.3 Quasy steady-state and quasiequilibrium asymptotics

Quasiequilibrium approximation uses the assumption that a group of reactions is much faster than other and goes fast to its equilibrium. We use below superscripts ‘f’ and ‘s’ to distinguish fast and slow reactions. A small parameter appears in the following form

$$\frac{dc}{dt} = \sum_{s, \text{ slow}} w_s^s(c, T) \gamma_s^s + \frac{1}{\varepsilon} \sum_{\varsigma, \text{ fast}} w_{\varsigma}^f(c, T) \gamma_{\varsigma}^f, \quad (4)$$

To separate variables, we have to study the spaces of linear conservation law of the initial system (4) and of the fast subsystem

$$\frac{dc}{dt} = \frac{1}{\varepsilon} \sum_{\varsigma, \text{ fast}} w_{\varsigma}^f(c, T) \gamma_{\varsigma}^f$$

If they coincide, then the fast subsystem just dominates, and there is no fast-slow separation for variables (all variables are either fast or constant). But if there exist additional linearly independent linear conservation laws for the fast system, then let us introduce new variables: linear functions $b^1(c), \dots, b^m(c)$, where $b^1(c), \dots, b^m(c)$ is the basis of the linear conservation laws for the initial system, and $b^1(c), \dots, b^{m+l}(c)$ is the basis of the linear conservation laws for the fast subsystem. Then

$b^{m+l+1}(c), \dots, b^n(c)$ are fast variables, $b^{m+1}(c), \dots, b^{m+l}(c)$ are slow variables, and $b^1(c), \dots, b^m(c)$ are constant. The *quasiequilibrium manifold* is given by the equations $\sum_{\zeta} w_{\zeta}^f(c, T) \gamma_{\zeta}^f = 0$ and for small ε it serves as an approximation to a slow manifold.

The quasi steady-state (or pseudo steady state) assumption was invented in chemistry for description of systems with radicals or catalysts. In the most usual version the species are split in two groups with concentration vectors c^s (“slow” or basic components) and c^f (“fast intermediates”). For catalytic reactions there is additional balance for c^f , amount of catalyst, usually it is just a sum $b_f = \sum_i c_i^f$. The amount of the fast intermediates is assumed much smaller than the amount of the basic components, but the reaction rates are of the same order, or even the same (both intermediates and slow components participate in the same reactions). This is the source of a small parameter in the system. Let us scale the concentrations c^f and c^s to the compatible amounts. After that, the fast and slow time appear and we could write $\dot{c}^s = W^s(c^s, c^f)$, $\dot{c}^f = \frac{1}{\varepsilon} W^f(c^s, c^f)$, where ε is small parameter, and functions W^s, W^f are bounded and have bounded derivatives (are “of the same order”). We can apply the standard singular perturbation techniques. If dynamics of fast components under given values of slow concentrations is stable, then the slow attractive manifold exists, and its zero approximation is given by the system of equations $W^f(c^s, c^f) = 0$.

The QE approximation is also extremely popular and useful. It has simpler dynamical properties (respects thermodynamics, for example, and gives no critical effects in fast subsystems of closed systems). Nevertheless, neither radicals in combustion, nor intermediates in catalytic kinetics are, in general, close to quasiequilibrium. They are just present in much smaller amount, and when this amount grows, then the QSS approximation fails.

The simplest demonstration of these two approximation gives the simple reaction: $S + E \leftrightarrow SE \rightarrow P + E$ with reaction rate constants $k_1^{\pm}$ and k_2 . The only possible quasiequilibrium appears when the first equilibrium is fast: $k_1^{\pm} = \kappa^{\pm}/\varepsilon$. The corresponding slow variable is $C^s = c_S + c_{SE}$, $b_E = c_E + c_{SE} = \text{const}$. For the QE manifold we get a quadratic equation $\frac{k_1^-}{k_1^+} c_{SE} = c_S c_E = (C^s - c_{SE})(b_E - c_{SE})$. This equation gives the explicit dependence $c_{SE}(C^s)$, and the slow equation reads $\dot{C}^s = -k_2 c_{SE}(C^s)$, $C^s + c_P = b_S = \text{const}$.

For the QSS approximation of this reaction kinetics, under assumption $b_E \ll b_S$, we have fast intermediates E and SE . For the QSS manifold there is a linear equation $k_1^+ c_S c_E - k_1^- c_{SE} - k_2 c_{SE} = 0$, which gives us the explicit expression for $c_{SE}(c_S)$: $c_{SE} = k_1^+ c_S b_E / (k_1^+ c_S + k_1^- + k_2)$ (the standard Michaelis–Menten formula). The slow kinetics reads $\dot{c}_S = -k_1^+ c_S (b_E - c_{SE}(c_S)) + k_1^- c_{SE}(c_S)$. The difference between the QSS and the QE in this example is obvious.

The terminology is not rigorous, and often QSS is used for all singular perturbed systems, and QE is applied only for the thermodynamic exclusion of fast variables by the maximum entropy (or minimum of free energy, or extremum of another relevant thermodynamic function) principle (MaxEnt). This terminological convention may be convenient. Nevertheless, without any relation to terminology, the difference between these two types of introduction of a small parameter is huge. There exists plenty of generalizations of these approaches, which aim to construct a slow and (almost) invariant manifold, and to approximate fast motion as well. The following references can give a first impression about these methods: Method of Invariant Manifolds (MIM) ([14, 46], Method of Invariant Grids (MIG), a discrete analogue of invariant manifolds ([13]), Computational Singular Perturbations (CSP) ([31, 32, 57]) Intrinsic Low-Dimensional Manifolds (ILDm) by [34], developed further in series of works by [4]), methods based on the Lyapunov auxiliary theorem ([24]).

2.4 Multiscale monomolecular reaction networks

A *monomolecular reaction* is such that it has at most one reactant and at most one product. Let us consider a general network of monomolecular reactions. This network is represented as a directed graph (digraph) [47]: vertices correspond to components A_i , edges correspond to reactions $A_i \rightarrow A_j$ with kinetic constants $k_{ji} > 0$. For each vertex, A_i , a positive real variable c_i (concentration) is defined. “Pseudo-species” (labeled $\emptyset$) can be defined to collect all degraded products, and degradation reactions can be written as $A_i \rightarrow \emptyset$ with constants k_{0i} . Production reactions can be represented as $\emptyset \rightarrow A_i$ with rates k_{i0} . The kinetic equation for the system is

$$\frac{dc_i}{dt} = k_{i0} + \sum_{j \geq 1} k_{ij} c_j - \sum_{j \geq 0} k_{ji} c_i, \quad (5)$$

or in vector form: $\dot{c} = K_0 + Kc$. Solution of this system can be reduced to a linear algebra problem: let us find all left (l^i) and right (r^i) eigenvectors of K , i.e.:

$$Kr^i = \lambda_i r^i, l^i K = \lambda_i l^i, \quad (6)$$

with the normalisation $(l^i, r^i) = \delta_{ij}$, where δ_{ij} is Kronecker's delta. Then the solution of (5) is

$$c(t) = c^s + \sum_{k=1}^n r^k (l^k, c(0) - c^s) \exp(-\lambda_k t), \quad (7)$$

where c^s is the steady state of the system (5), i.e. when all $\frac{dc_i}{dt} = 0$, and $c(0)$ is the initial condition.

If all reaction constants k_{ij} would be known with precision then the eigenvalues and the eigenvectors of the kinetic matrix can be easily calculated by standard numerical techniques. Furthermore, Singular Value Decomposition (SVD) can be used for model reduction. But in systems biology models often one has only approximate or relative values of the constants (information on which constant is bigger or smaller than another one). Let us consider the simplest case: when all kinetic constants are very different (separated), i.e. for any two different pairs of indices $I = (i, j)$, $J = (i', j')$ we have either $k_I \gg k_J$ or $k_J \ll k_I$. In this case we say that the system is hierarchical with timescales (inverses of constants k_{ij} , $j \neq 0$) totally separated.

Linear network with totally separated constants can be represented as a digraph and a set of orders (integer numbers) associated to each arc (reaction). The lower the order, the more rapid is the reaction. It happens that in this case the special structure of the matrix K (originated from a reaction graph) allows us to exploit the strong relation between the dynamics (5) and the topological properties of the digraph. In this case, possible values of l_i are 0, 1 and the possible values of r_i are -1, 0, 1 with high precision. In previous works, we provided an algorithm for finding non-zero components of l_i , r_i , based on the network topology and the constants ordering, which gives a good approximation to the problem solution [16, 17, 45].

2.5 Dominant system for a simple irreversible catalytic cycle with limiting step

A linear chain of reactions, $A_1 \rightarrow A_2 \rightarrow \dots A_n$, with reaction rate constants k_i (for $A_i \rightarrow A_{i+1}$), gives the first example of limiting steps. Let the reaction rate constant k_q be the smallest one. Then we expect the following behaviour of the reaction chain in time scale $\gtrsim 1/k_q$: all the components $A_1, \dots A_{q-1}$ transform fast into A_q , and all the components $A_{q+1}, \dots A_{n-1}$ transform fast into A_n , only two components, A_q and A_n are present (concentrations of other components are small), and the whole dynamics in this time scale can be represented by a single reaction $A_q \rightarrow A_n$ with reaction rate constant k_q . This picture becomes more exact when k_q becomes smaller with respect to other constants.

The kinetic equation for the linear chain is


$$\dot{c}_i = k_{i-1} c_{i-1} - k_i c_i, \quad (8)$$

The coefficient matrix K of these equations is very simple. It has nonzero elements only on the main diagonal, and one position below. The eigenvalues of K are $-k_i$ ($i = 1, \dots, n-1$) and 0. The left and right eigenvectors for 0 eigenvalue, l^0 and r^0 , are:

$$l^0 = (1, 1, \dots, 1), \quad r^0 = (0, 0, \dots, 0, 1), \quad (9)$$

all coordinates of l^0 are equal to 1, the only nonzero coordinate of r^0 is r_n^0 and we represent vector-column r^0 in row.

The catalytic cycle is one of the most important substructures that we study in reaction networks. In the reduced form the catalytic cycle is a set of linear reactions:



Reduced form means that in reality some of these reaction are not monomolecular and include some other components (not from the list $A_1, \dots A_n$). But in the study of the isolated cycle dynamics,

concentrations of these components are taken as constant and are included into kinetic constants of the cycle linear reactions.

For the constant of elementary reaction $A_i \rightarrow$ we use the simplified notation k_i because the product of this elementary reaction is known, it is A_{i+1} for $i < n$ and A_1 for $i = n$. The elementary reaction rate is $w_i = k_i c_i$, where c_i is the concentration of A_i . The kinetic equation is:

$$\dot{c}_i = k_{i-1} c_{i-1} - k_i c_i, \quad (10)$$

where by definition $c_0 = c_n$, $k_0 = k_n$, and $w_0 = w_n$. In the stationary state ($\dot{c}_i = 0$), all the w_i are equal: $w_i = w$. This common rate w we call the cycle stationary rate, and

$$w = \frac{b}{\frac{1}{k_1} + \dots + \frac{1}{k_n}}; \quad c_i = \frac{w}{k_i}, \quad (11)$$

where $b = \sum_i c_i$ is the conserved quantity for reactions in constant volume. Let one of the constants, $k_{\min}$, be much smaller than others (let it be $k_{\min} = k_n$):

$$k_i \gg k_{\min} \quad \text{if } i \neq n. \quad (12)$$

In this case, in linear approximation

$$c_n = b \left(1 - \sum_{i < n} \frac{k_n}{k_i} \right), \quad c_i = b \frac{k_n}{k_i}, \quad w = k_n b. \quad (13)$$

The simplest zero order approximation for the steady state gives

$$c_n = b, \quad c_i = 0 \quad (i \neq n). \quad (14)$$

This is trivial: all the concentration is collected at the starting point of the “narrow place”, but may be useful as an origin point for various approximation procedures.

So, the stationary rate of a cycle is determined by the smallest constant, $k_{\min}$, if it is much smaller than the constants of all other reactions (12):

$$w \approx k_{\min} b. \quad (15)$$

In that case we say that the cycle has a limiting step with constant $k_{\min}$.

There is significant difference between the examples of limiting steps for the chain of reactions and for irreversible cycle. For the chain, the steady state does not depend on nonzero rate constants. It is just $c_n = b, c_1 = c_2 = \dots = c_{n-1} = 0$. The smallest rate constant k_q gives the smallest positive eigenvalue, the relaxation time is $\tau = 1/k_q$. The corresponding approximation of eigenmode (right eigenvector) r^1 has coordinates: $r_1^1 = \dots = r_{q-1}^1 = 0, r_q^1 = 1, r_{q+1}^1 = \dots = r_{n-1}^1 = 0, r_n^1 = -1$. This exactly corresponds to the statement that the whole dynamics in the time scale $\gtrsim 1/k_q$ can be represented by a single reaction $A_q \rightarrow A_n$ with reaction rate constant k_q . The left eigenvector for eigenvalue k_q has approximation l^1 with coordinates $l_1^1 = l_2^1 = \dots = l_q^1 = 1, l_{q+1}^1 = \dots = l_n^1 = 0$. This vector provides the almost exact *lumping* on time scale $\gtrsim 1/k_q$. Let us introduce a new variable $c_{\text{lump}} = \sum_i l_i c_i$, i.e. $c_{\text{lump}} = c_1 + c_2 + \dots + c_q$. For the time scale $\gtrsim 1/k_q$ we can write $c_{\text{lump}} + c_n \approx b$, $dc_{\text{lump}}/dt \approx -k_q c_{\text{lump}}, dc_n/dt \approx k_q c_{\text{lump}}$.

In the example of a cycle, we approximate the steady state, that is, the right eigenvector r^0 for zero eigenvalue (the left eigenvector is known and corresponds to the main linear balance b : $l_i^0 \equiv 1$). In the zero-order approximation, this eigenvector has coordinates $r_1^0 = \dots = r_{n-1}^0 = 0, r_n^0 = 1$.

If k_n/k_i is small for all $i < n$, then the kinetic behaviour of the cycle is determined by a linear chain of $n-1$ reactions $A_1 \rightarrow A_2 \rightarrow \dots A_n$, which we obtain after cutting the limiting step. The characteristic equation for an irreversible cycle, $\prod_{i=1}^n (\lambda + k_i) - \prod_{i=1}^n k_i = 0$, tends to the characteristic equation for the linear chain, $\lambda \prod_{i=1}^{n-1} (\lambda + k_i) = 0$, when $k_n \rightarrow 0$.

The characteristic equation for a cycle with limiting step ($k_n/k_i \ll 1$) has one simple zero eigenvalue that corresponds to the conservation law $\sum c_i = b$ and $n-1$ nonzero eigenvalues

$$\lambda_i = -k_i + \delta_i \quad (i < n). \quad (16)$$

where $\delta_i \rightarrow 0$ when $\sum_{i < n} \frac{k_n}{k_i} \rightarrow 0$.

A cycle with limiting step (10) has real eigenspectrum and demonstrates monotonic relaxation without damped oscillations. Of course, without limitation such oscillations could exist, for example, when all $k_i \equiv k > 0$, ($i = 1, \dots, n$).

The relaxation time of a stable linear system (10) is, by definition, $\tau = 1/\min\{Re(-\lambda_i)\}$ ($\lambda \neq 0$). For small k_n , $\tau \approx 1/k_\tau$, $k_\tau = \min\{k_i\}$, ($i = 1, \dots, n-1$). In other words, for a cycle with limiting step, k_τ is the *second slowest rate constant*: $k_{\min} \ll k_\tau \leq \dots$

3 Results

3.1 Model assumptions

We consider two models of action of microRNA on protein translation process proposed in [37]: the simplest linear model, and the non-linear model which explicitly takes into account recycling of ribosomal subunits and initiation factors.

Both models, of course, represent significant simplifications of biological reality. First, only a limited subset of all possible mechanisms of microRNA action on the translation process is considered (see Fig. 1). Second, all processes of synthesis and degradation of mRNA and protein are deliberately neglected. Third, interaction of microRNA and mRNA is simplified: it is supposed that when microRNA is added to the experimental system then only mRNA with bound microRNAs are present (this also assumes that the concentration of microRNA is abundant with respect to mRNA). Concentrations of microRNA and mRNA are supposed to be constant. Interaction of only one type of microRNA and one type of mRNA is considered (not a mix of several microRNAs). The process of initiation is greatly simplified: all initiation factors are represented by only one molecule which is marked as eIF4F.

Finally, the classical chemical kinetics approach is applied, based on solutions of ordinary differential equations, which supposes sufficient and well-stirred amount of both microRNAs and mRNAs. Another assumption in the modeling is the mass action law assumed for the reaction kinetic rates.

It is important to underline the interpretation of certain chemical species considered in the system. The ribosomal subunits and the initiation factors in the model exist in free and bound forms, moreover, ribosomal subunits can be bound to several regions of mRNA (the initiation site, the start codon, the coding part). Importantly, several copies of fully assembled ribosome can be bound to one mRNA. To model this situation, we have to introduce the following quantification rule for chemical species: amount of "ribosome bound to mRNA" means the total number of ribosomes translating proteins, which is not equal to the number of mRNAs with ribosome sitting on them, since one mRNA can hold several translating ribosomes (polyribosome). In this view, mRNAs act as *places* or *catalyzers*, where translation takes place, whereas mRNA itself formally is not consumed in the process of translation, but, of course, can be degraded or synthesized (which is, however, not considered in the models described further).

3.2 The simplest linear protein translation model

The simplest representation of the translation process has the form of a circular cascade of reactions [37] (see Fig. 2).

The list of chemical species in the model is the following:

1. 40S, free small ribosomal subunit.
2. mRNA:40S, small ribosomal subunit bound to the initiation site.
3. AUG, small ribosomal subunit bound to the start codon.

The catalytic cycle is formed by the following reactions:

1. $40S \rightarrow \text{mRNA:40S}$, Initiation complex assembly (rate k_1).
2. $\text{mRNA:40S} \rightarrow \text{AUG}$, Some late and cap-independent initiation steps, such as scanning the 5'UTR by for the start codon AUG recognition (rate k_2).
3. $\text{AUG} \rightarrow 40S$, combined processes of 60S ribosomal unit joining and protein elongation, which leads to production of the protein (rate k_3), and fall off of the ribosome from mRNA.

The model is described by the following system of equations [37]:

$$\begin{cases} \frac{d[40S](t)}{dt} = -k_1[40S] + k_3[AUG] \\ \frac{d[mRNA : 40S](t)}{dt} = k_1[40S] - k_2[mRNA : 40S] \\ \frac{d[AUG](t)}{dt} = k_2[mRNA : 40S] - k_3[AUG] \\ Prsynth(t) = k_3[AUG](t) \end{cases} \quad (17)$$

where $Prsynth(t)$ is the rate of the protein synthesis.

Following [37], let us assume that $k_3 \gg k_1, k_2$. This choice was justified by the following statement: “...The subunit joining and protein production rate (k_3) is faster than k_1 and k_2 since mRNA40S complexes bound to the AUG without the 60S subunit are generally not observed in translation initiation unless this step is stalled by experimental methods, and elongation is generally thought to not be rate limiting in protein synthesis.” [37].

Under this condition, the equations (17) have the following approximate solution (which becomes the more exact the smaller the $\frac{k_1+k_2}{k_3}$ ratio):

$$\begin{bmatrix} [40S](t) \\ [mRNA : 40S](t) \\ [AUG](t) \end{bmatrix} = \frac{[40S]_0}{\frac{1}{k_1} + \frac{1}{k_2}} \left(\begin{bmatrix} 1/k_1 \\ 1/k_2 \\ 1/k_3 \end{bmatrix} + \frac{1}{k_3} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} e^{-k_3 t} + \frac{1}{k_2} \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-(k_1+k_2)t} \right) \quad (18)$$

$$Prsynth(t) = \frac{[40S]_0}{\frac{1}{k_1} + \frac{1}{k_2}} \left(1 - \frac{k_3}{k_2} e^{-(k_1+k_2)t} \right) \quad (19)$$

for the initial condition

$$\begin{bmatrix} [40S] \\ [mRNA : 40S] \\ [AUG] \\ Prsynth \end{bmatrix}_0 = \begin{bmatrix} [40S]_0 \\ 0 \\ 0 \\ 0 \end{bmatrix}. \quad (20)$$

From the solution (18) it follows that the dynamics of the system evolves on two time scales: 1) fast elongation dynamics on the time scale $\approx 1/k_3$; and 2) relatively slow translation initiation dynamics with the relaxation time $t_{rel} \approx \frac{1}{k_1+k_2}$. The protein synthesis rate formula (19) does not include the k_3 rate, since it is neglected with respect to k_1, k_2 values. From (19) we can extract the formula for the protein synthesis steady-state rate $Prsynth$ (multiplier before the parentheses) and the relaxation time t_{rel} for it (inverse of the exponent power):

$$Prsynth = \frac{[40S]_0}{\frac{1}{k_1} + \frac{1}{k_2}}, \quad t_{rel} = \frac{1}{k_1 + k_2} \quad (21)$$

Now let us consider two experimental situations: 1) the rates of the two translation initiation steps are comparable $k_1 \approx k_2$; 2) the cap-dependent rate k_1 is limiting: $k_1 \ll k_2$. Accordingly to [37], the second situation can correspond to modified mRNA with an alternative cap-structure, which is much less efficient for the assembly of the initiation factors, 40S ribosomal subunit and polyA binding proteins.

For these two experimental systems (let us call them ‘wild-type’ and ‘modified’ correspondingly), let us study the effect of microRNA action. We will model the microRNA action as a mechanism which effectively diminish the value of a kinetic rate coefficient. Let us assume that there are two alternative mechanisms: 1) microRNA acts in a cap-dependent manner (thus, reducing the k_1 constant) and 2) microRNA acts in a cap-independent manner, for example, through interfering with 60S subunit joining (thus, reducing the k_2 constant). The dependence of the steady rate of protein synthesis $Prsynth \sim \frac{1}{\frac{1}{k_1} + \frac{1}{k_2}}$ and the relaxation time $t_{rel} \approx \frac{1}{k_1+k_2}$ on the efficiency of the microRNA action (i.e., how much it is capable to diminish a rate coefficient) is shown on Fig. 3.

Interestingly, experiments with cap structure replacement were made and the effect of microRNA action on the translation was measured [36, 48]. No change in the protein rate synthesis after applying




Figure 2: The simplest model of microRNA action on the protein translation, represented with use of Systems Biology Graphical Notation (a) and schematically with the condition on the constants (b). The two mechanism of microRNA action.

microRNA was observed. From this it was concluded that microRNA in this system should act through cap-dependent mechanism (i.e., the normal 'wild-type' cap is required for microRNA recruitment). It was argued that this could be a misinterpretation [37] since in the 'modified' system, cap-dependent translation initiation is a rate limiting process ($k_1 \ll k_2$), hence, even if microRNA acts in the cap-independent manner (inhibiting k_2), it will have no effect on the final steady state protein synthesis rate. They confirmed this by the graph similar to the Fig. 3a.

From the analytical solution (18) we can further develop this idea and claim that it is still possible to detect the action of microRNA in the 'modified' system if one measures the protein synthesis relaxation time: if it significantly increases then microRNA probably acts in the cap-independent manner despite the fact that the steady state rate of the protein synthesis does not change (see the Fig. 3b). This is a simple consequence of the fact that the relaxation time in a cycle of biochemical reactions is limited by the second slowest reaction [17]. If the relaxation time is not changed in the presence of microRNA then we can conclude that none of the two alternative mechanisms of microRNA-based translation repression is activated in the system, hence, microRNA action is dependent on the structure of the 'wild-type' transcript cap.

The observations from the Fig. 3 are recapitulated in the Table 1. This analysis (of course, oversimplified in many aspects) provides us with an important lesson: observed dynamical features of the translation process with and without presence of microRNA can give clues on the mechanisms of microRNA action and help to distinguish them in a particular experimental situation. Theoretical analysis of the translation dynamics highlights what are the important characteristics of the dynamics which should be measured in order to infer the possible microRNA mechanism.

This conclusion suggests the notion of a *kinetic signature of microRNA action mechanism* which we define as *the set of measurable characteristics of the translational machinery dynamics (features of time series for protein, mRNA, ribosomal subunits concentrations) and the predicted tendencies of their changes as a response to microRNA action through a particular biochemical mechanism*.

3.3 The non-linear protein translation model

To explain the effect of microRNA interference with translation initiation factors, a non-linear version of the translation model was proposed [37] which explicitly takes into account recycling of initiation factors (eIF4F) and ribosomal subunits (40S and 60S).

The model contains the following list of chemical species (see also Fig. 4):

1. 40S, free 40S ribosomal subunit.
2. 60S, free 60S ribosomal subunit.
3. eIF4F, free initiation factor.




Figure 3: Graphs illustrating the predicted change in the steady-state rate of protein synthesis (left), and its relaxation time (i.e., the time needed to recover from a perturbation to the steady state value), right. Four curves are presented. The black ones are for the wild-type cap structure, which is modeled by $k_1 = k_2$. The red ones are for the modified structure, when $k_1 \ll k_2$. The main conclusion from the left graph is that if microRNA acts on a late initiation step, diminishing k_2 then it's effect is not measurable unless k_2 is very strongly suppressed (as reported in [37]). The main conclusion from the right is that the effect of microRNA should be measurable in this case if one looks at dynamical features such the relaxation time.

Table 1: Modeling of two mechanisms of microRNA action in the simplest linear model. microRNA action effect is described for the protein synthesis steady rate and the relaxation time. It is assumed that the ribosome assembly+elongation step in protein translation, described by the k_3 rate constant, is not rate limiting

Observable value	Initiation(k_1)	Step after initiation, cap-independent(k_2)	Elongation (k_3)
Wild-type cap			
<i>Steady-state rate</i>	decreases	decreases	no change
<i>Relaxation time</i>	increases slightly	increases slightly	no change
A-cap			
<i>Steady-state rate</i>	decreases	no change	no change
<i>Relaxation time</i>	no change	increases drastically	no change

4. mRNA:40S, formed initiation complex (containing 40S and the initiation factors), bound to the initiation site of mRNA.

5. AUG, initiation complex bound to the start codon of mRNA.

6. 80S, fully assembled ribosome translating protein.

There are four reactions in the model, all considered to be irreversible:

1. $40S + eIF4F \rightarrow \text{mRNA:40S}$, assembly of the initiation complex (rate k_1).

2. $\text{mRNA:40S} \rightarrow \text{AUG}$, some late and cap-independent initiation steps, such as scanning the 5'UTR by for the start codon AUG recognition (rate k_2).

3. $\text{AUG} \rightarrow 80S$, assembly of ribosomes and protein translation (rate k_3).

4. $80S \rightarrow 60S + 40S$, recycling of ribosomal subunits (rate k_4).

The model is described by the following system of equations [37] :

$$\begin{cases} \frac{d[40S]}{dt} = -k_1[40S][eIF4F] + k_4[80S] \\ \frac{d[eIF4F]}{dt} = -k_1[40S][eIF4F] + k_2[\text{mRNA:40S}] \\ \frac{d[\text{mRNA:40S}]}{dt} = k_1[40S][eIF4F] - k_2[\text{mRNA:40S}] \\ \frac{d[AUG]}{dt} = k_2[\text{mRNA:40S}] - k_3[AUG][60S] \\ \frac{d[60S]}{dt} = -k_3[AUG][60S] + k_4[80S] \\ \frac{d[80S]}{dt} = k_3[AUG][60S] - k_4[80S] \\ Prsynth(t) = k_3[AUG][60S] \end{cases} \quad (22)$$

where $[40S]$ and $[60S]$ are the concentrations of free 40S and 60S ribosomal subunits, $[eIF4F]$ is a concentration of free translation initiation factors, $[\text{mRNA:40S}]$ is the concentration of 40S subunit bound to the initiation site of mRNA, $[AUG]$ is the concentration of the initiation complex bound to the start codon, $[80S]$ is the concentration of ribosomes translating protein, and $Prsynth$ is the rate of protein synthesis.

The model (22) contains three independent conservations laws:

$$[\text{mRNA:40S}] + [40S] + [AUG] + [80S] = [40S]_0, \quad (23)$$

$$[\text{mRNA:40S}] + [eIF4F] = [eIF4F]_0, \quad (24)$$

$$[60S] + [80S] = [60S]_0, \quad (25)$$

The following assumptions on the model parameters were suggested in [37]:

$$k_4 \ll k_1, k_2, k_3; k_3 \gg k_1, k_2; [eIF4F]_0 \ll [40S]_0; [eIF4F]_0 < [60S]_0 < [40S]_0 \quad (26)$$

with the following justification: "...The amount 40S ribosomal subunit was set arbitrarily high ... as it is thought to generally not be a limiting factor for translation initiation. In contrast, the level of eIF4F, as the canonical limiting factor, was set significantly lower so translation would be dependent on its concentration as observed experimentally... Finally, the amount of subunit joining factors for the 60S large ribosomal subunit were estimated to be more abundant than eIF4F but still substoichiometric when compared to 40S levels, consistent with in vivo levels... The k_4 rate is relatively slower than the other rates in the model; nevertheless, the simulations overall protein production was not altered by changes of several orders of magnitude around its value... " [37].

Notice that further in our paper we show that the last statement about the value of k_4 is needed to be made more precise: in the model by Nissan and Parker, k_4 is a critical parameter. It does not affect the steady state protein synthesis rate only in one of the possible scenarios (inefficient initiation, deficit of the initiation factors).

3.3.1 Steady state solution

The final steady state of the system can be calculated from the conservation laws and the balance equations among all the reaction fluxes:

$$k_2 \cdot [mRNA : 40S]_s = k_3 \cdot [AUG]_s \cdot [60S]_s = k_4 \cdot [80S]_s = k_1 \cdot [40S]_s \cdot [eIF4F]_s, \quad (27)$$

where 's' index stands for the steady state value. Let us designate a fraction of the free [60S] ribosomal subunit in the steady state as $x = \frac{[60S]_s}{[60S]_0}$. Then we have

$$\begin{aligned} [mRNA : 40S]_s &= \frac{k_4}{k_2} [60S]_0 (1 - x), [AUG]_s = \frac{k_4}{k_3} \left(\frac{1 - x}{x} \right), \\ [eIF4F]_s &= [eIF4F]_0 - \frac{k_4}{k_2} [60S]_0 (1 - x), \\ [60S]_s &= [60S]_0 x, [80S]_s = [60S]_0 (1 - x), \\ [40S]_s &= [40S]_0 - [60S]_0 (1 - x) \left(1 + \frac{k_4}{k_2} \right) - \frac{k_4}{k_3} \left(\frac{1 - x}{x} \right) \end{aligned} \quad (28)$$

and the equation to determine x , in which we have neglected the terms of smaller order of magnitude, based on conditions (26):

$$\begin{aligned} x^3 + x^2 (\alpha + (\delta - 1) + (\beta - 1)) + x (-\alpha + (\delta - 1)(\beta - 1)) + \gamma(1 - \beta) &= 0, \\ \alpha = \frac{k_2}{k_1 [60S]_0}, \beta = \frac{[eIF4F]_0 k_2}{[60S]_0 k_4}, \gamma = \frac{k_4}{k_3 [60S]_0}, \delta = \frac{[40S]_0}{[60S]_0} \end{aligned} \quad (29)$$

From the inequalities on the parameters of the model, we have $\delta > 1$, $\gamma \ll 1$ and, if $k_1 \gg k_4/[eIF4F]_0$ then $\alpha \ll \beta$. From these remarks it follows that the constant term $\gamma(1 - \beta)$ of the equation (29) should be much smaller than the other polynomial coefficients, and the equation (29) should have one solution close to zero and two others:

$$\begin{aligned} x_0 &\approx \frac{k_4}{k_3 \cdot ([40S]_0 - [60S]_0)}, \\ x_1 &\approx 1 - \beta + \frac{\alpha\beta}{\delta - \beta} = 1 - \frac{[eIF4F]_0 k_2}{[60S]_0 k_4} + \frac{k_2^2 [eIF4F]_0}{k_1 k_4 [40S]_0} \frac{1}{1 - \frac{[eIF4F]_0 k_2}{[40S]_0 k_4}}, \\ x_2 &\approx 1 - \frac{[40S]_0}{[60S]_0}, \end{aligned} \quad (30)$$

provided that $\alpha \ll |1 - \delta|$ or $\alpha \ll |1 - \beta|$. In the expression for x_1 we can not neglect the term proportional to α , to avoid zero values in (29).

The solution x_2 is always negative, which means that one can have one positive solution $x_0 \ll 1$ if $\frac{[eIF4F]_0 k_2}{[60S]_0 k_4} \gtrsim 1$ and two positive solutions x_0 and x_1 if $\frac{[eIF4F]_0 k_2}{[60S]_0 k_4} \lesssim 1$. However, from (28), (30) and (26) it is easy to check that if $x_1 > 0$ then x_0 does not correspond to positive value of $[eIF4F]_s$. This means that for a given combination of parameters satisfying (26) we can have only one steady state (either x_0 or x_1).

The two values $x = x_0$ and $x = x_1$ correspond to two different modes of translation. When, for example, the amount of the initiation factors $[eIF4F]_0$ is not sufficient to provide efficient initiation ($[eIF4F]_0 < \frac{k_2}{[60S]_0 k_4}$, $x = x_1$) then most of the 40S and [60S] subunits remain in the free form, the initiation factor $[eIF4F]$ being always the limiting factor. If the initiation is efficient enough ($[eIF4F]_0 > \frac{k_2}{[60S]_0 k_4}$), then we have $x = x_0 \ll 1$ when almost all 60S ribosomal subunits are engaged in the protein elongation, and $[eIF4F]$ being a limiting factor at the early stage, however, is liberated after and ribosomal subunits recycling becomes limiting in the initiation (see the next section for the analysis of the dynamics).

Let us notice that the steady state protein synthesis rate under these assumptions is




Figure 4: Non-linear model of microRNA action on the protein translation, represented with use of Systems Biology Graphical Notation (a) and schematically with the condition on the constants (b). The difference from the simplest model (2) is in the explicit description of initiation factors $eIF4F$, and ribosomal subunits 40S and 60S recycling.

$$Prsynth = k_4 \cdot [60S]_0(1 - x) \approx \begin{cases} k_4 \cdot [60S]_0, & \text{if } \frac{[eIF4F]_0 k_2}{[60S]_0 k_4} > 1 \\ k_2 \cdot [eIF4F]_0, & \text{else} \end{cases} \quad (31)$$

This explains the numerical results obtained in [37]: with low concentrations of $[eIF4F]_0$ microRNA action would be efficient only if it affects k_2 or if it competes with $eIF4F$ for binding to the mRNA cap structure (thus, effectively further reducing the level $[eIF4F]_0$). With higher concentrations $[eIF4F]_0$, other limiting factors become dominant: $[60S]_0$ (availability of the heavy ribosomal subunit) and k_4 (speed of ribosomal subunits recycling which is the slowest reaction rate in the system). Interestingly, in any situation the protein translation rate does not depend on the value of k_1 directly (of course, unless it does not become "globally" rate limiting), but only through competing with $eIF4F$ (which makes the difference with the simplest linear protein translation model).

Equation (31) explains also some experimental results reported in [36]: increasing the concentration of $[eIF4F]$ translation initiation factor enhances protein synthesis but its effect is abruptly saturated above a certain level.

It would be interesting to make some conclusions on the shift of polysomal profile from the steady state solutions 30. In this model, the number of ribosomes sitting on mRNA $N_{polysome}$ is defined by $N_{polysome} = \frac{[80S]}{[mRNA]}$, where $[mRNA]$ is the concentration of mRNA. However, $[mRNA]$ is not an explicit dynamical variable in the model, it is implicitly included in other model constants, such as k_1 , together with the effective volume of cytoplasmic space considered in the model. Nevertheless, the model can predict the relative shift of the polysome profile. In the steady state

$$N_{polysome} \sim 1 - x = \begin{cases} k_4 \cdot [60S]_0, & \text{if } \frac{[eIF4F]_0 k_2}{[60S]_0 k_4} > 1 \\ k_2 \cdot [eIF4F]_0, & \text{else} \end{cases} \quad (32)$$

and $N_{polysome}$ changes in the same way as the protein synthesis steady state value.

3.3.2 Analysis of the dynamics

It was proposed to use the following model parameters in [37]: $k_1 = k_2 = 2$, $k_3 = 5$, $k_4 = 1$, $[40S]_0 = 100$, $[60S]_0 = 25$, $[eIF4F]_0 = 6$. Although all estimations made in the previous section remain valid for them, in this section we have modified the parameter values (anyway very arbitrary estimated) to make the dynamical features of the model more illustrative. After we have verified that the semi-analytical approach remains valid for the set of "normal" parameter values from [37] (and, in general, for a very large set of parameters). First, we modified $k_2 = 3$ to avoid possible artifact symmetries related to equal parameter values. Second, we have changed $k_4 = 0.1$ to have the case of $x = x_0$ steady state (which, as it will be shown, corresponds to the three-stage dynamics, which allows

to demonstrate three dominant dynamical systems instead of only one in the case of $x = x_1$ when $eIF4F$ is a limiting factor during the whole relaxation process) and $k_3 = 50$ to have better separation between k_3 and k_1, k_2 .

Simulations of the protein translation model with these parameters and the initial conditions

$$\begin{bmatrix} [40S] \\ [eIF4F] \\ [mRNA : 40S] \\ [AUG] \\ [80S] \\ [60S] \end{bmatrix}_0 = \begin{bmatrix} [40S]_0 \\ [eIF4F]_0 \\ 0 \\ 0 \\ 0 \\ [60S]_0 \end{bmatrix}. \quad (33)$$

are shown on the Fig. 6. The system shows non-trivial relaxation process which takes place in several epochs. One can check that the "efficient translation initiation" ($x = x_0 \ll 1$) scenario is realized in this case. Qualitatively we can distinguish the following stages:

1) Stage 1: Relatively fast relaxation with conditions $[40S] \gg [eIF4F]$, $[60S] \gg [AUG]$. During this stage, the two non-linear reactions $40S + eIF4F \rightarrow mRNA : 40S$ and $AUG + 60S \rightarrow 80S$ can be considered as pseudo-monomolecular ones: $eIF4F \rightarrow mRNA : 40S$ and $AUG \rightarrow 80S$ with rate constants dependent on $[40S]$ and $[60S]$ respectively. This stage is characterized by rapidly establishing quasiequilibrium of three first reactions (R1, R2 and R3 with k_1, k_2 and k_3 constants). Biologically, this stage corresponds to assembling of the translation initiation machinery, scanning for the start codon and assembly of the first full ribosome at the start codon position.

2) Transition between Stage 1 and Stage 2.

3) Stage 2: Relaxation with the conditions $[40S] \gg [eIF4F]$, $[60S] \ll [AUG]$. During this stage, the reactions $40S + eIF4F \rightarrow mRNA : 40S$ and $AUG + 60S \rightarrow 80S$ can be considered as pseudo-monomolecular $eIF4F \rightarrow mRNA : 40S$ and $60S \rightarrow 80S$. This stage is characterized by two local quasi-steady states established in the two network reaction cycles (formed from R1-R2 and R3-R4 reactions). Biologically, this stage corresponds to the first round of elongation, when first ribosomes moves along the coding region of mRNA. The small ribosomal subunit 40S is still in excess which keeps the initiation stage (reaction R1-R2 fluxes) relatively fast.

4) Transition between Stage 2 and Stage 3.

5) Stage 3: Relaxation with the conditions $[40S] \ll [eIF4F]$, $[60S] \ll [AUG]$. During this stage, the reactions $40S + eIF4F \rightarrow mRNA : 40S$ and $AUG + 60S \rightarrow 80S$ can be considered as pseudo-monomolecular $40S \rightarrow mRNA : 40S$ and $60S \rightarrow 80S$. During this stage all reaction fluxes are balanced. Biologically, this stage corresponds to the stable production of the protein with constant recycling of the ribosomal subunits. Most of ribosomal subunits 40S are involved in protein elongation, so the initiation process should wait until the end of elongation for that they would be recycled.

Stages 1-3 can be associated with the corresponding dominant systems [17] which are shown on Fig. 5.

Below we give a more detailed analysis of stages 1-3 and transitions between them.

3.3.3 Stage 1: translation initiation and assembly of the first ribosome at the start codon

The dominant system of the Stage 1 (Fig. 5a) can be modeled as a linear system of equations (notice that it is not equivalent to the system of equations that would correspond to fully monomolecular reaction network because the reaction R2 is still bimolecular despite the fact that the products of this reaction do not interact, which leads to the linear description):




Figure 5: Dominant systems for three stages of relaxation of the model (22). Stage 1) The dominant system is pseudo-linear network of reactions. Stage 2) The dominant system is a quasi-steady state approximation, where one supposes that the fluxes in two network cycles are balanced. Stage 3) The dominant system is pseudo-linear network of reactions.

$$\begin{cases} \frac{d[eIF4F]}{dt} = -k'_1[eIF4F] + k_2[mRNA : 40S] \\ \frac{d[mRNA : 40S]}{dt} = k'_1[eIF4F] - k_2[mRNA : 40S] \\ \frac{d[AUG]}{dt} = k_2[mRNA : 40S] - k'_3[AUG] \\ \frac{d[80S]}{dt} = k_3[AUG] [60S] - k_4[80S] \end{cases} \quad (34)$$

where $k'_1 = k_1 \cdot [40S]$, $k'_3 = k_3 \cdot [60S]$ and we consider that at this stage the changes of 40S and 60S are relatively slow. This system has simple approximate solution, taking into account constraints on the parameters $k_2 \ll k'_1, k'_3$; $k_4 \ll k'_1, k'_3, k_2$, also assuming $k_2 \ll |k'_1 - k'_3|$, and for the initial condition

$$\begin{bmatrix} [eIF4F] \\ [mRNA : 40S] \\ [AUG] \\ [80S] \end{bmatrix}_0 = \begin{bmatrix} [eIF4F]_0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad (35)$$

we have

$$\begin{bmatrix} [eIF4F](t) \\ [mRNA : 40S](t) \\ [AUG](t) \\ [80S](t) \end{bmatrix} = k_2 \cdot [eIF4F]_0 \left(\begin{bmatrix} 1/k'_1 \\ 1/k_2 \\ 1/k'_3 \\ 1/k_4 \end{bmatrix} - \frac{1}{k'_3} \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix} e^{-k'_3 t} + \frac{1}{k_2} \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} e^{-k'_1 t} - \frac{1}{k_4} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} e^{-k_4 t} \right) \quad (36)$$

From this solution, one can conclude that the relaxation of this model goes at several time scales (very rapid $\sim \min(1/k'_3, 1/k'_1)$ and slow $\sim 1/k_4$) and that when eIF4F, mRNA:40S and AUG already reached their quasiequilibrium values, [80S] continues to grow. This corresponds to the quasiequilibrium approximation asymptotic (see section 2.3). At some point 80S will reach such a value that it would be not possible to consider 60S constant: otherwise the conservation law (25) will be violated. This will happen when $[80S] \ll [60S]$ condition is not satisfied anymore, i.e., following our convention to consider "much smaller" as difference in one order of magnitude, at $t' \sim \frac{[60S]_0}{10 \cdot [eIF4F]_0 \cdot k_2}$. The same consideration is applicable for another conservation law (23) in which [80S] is included, but from the time point $t'' \sim \frac{[40S]_0}{10 \cdot [eIF4F]_0 - 1}$. From $[40S]_0 \gg [eIF4F]_0$ and $[40S]_0 > [60S]_0$ we have $\min(t', t'') = t'$.




Figure 6: Simulation of the non-linear protein translation model with parameters $k_1 = 2$, $k_2 = 3$, $k_3 = 50$, $k_4 = 0.1$, $[40S]_0 = 100$, $[60S]_0 = 25$, $[eIF4F]_0 = 6$. a) and b) chemical species concentrations at logarithmic and linear scales; c) and d) reaction fluxes at logarithmic and linear scales. By the dashed line several stages are delimited during which the dynamics can be considered as (pseudo-)linear. To determine where " $\gg$ " and " $\ll$ " conditions are violated, we arbitrarily consider "much bigger" or "much smaller" as difference in one order of magnitude (by factor 10).

This means that the parameters $[40S]$, $[60S]$ of the (local) steady states for $[eIF4F]$ and $[AUG]$ should slowly (at the same rate as $[80S]$) change from the time point t' (variable $[mRNA : 40S]$ does not change because its local steady state does not depend on $[40S]$, $[60S]$). In other words, after $t = t'$ the Stage 1 solution (36) should be prolonged as

$$\begin{aligned} [80S](t) &= \frac{k_2 \cdot [eIF4F]_0}{k_4} (1 - e^{-k_4 t}), [40S](t) = [40S]_0 - [80S](t), [60S](t) = [60S]_0 - [80S](t), \\ [eIF4F](t) &= \frac{k_2 \cdot [eIF4F]_0}{k_1([40S]_0 - [80S](t))}, [mRNA : 40S](t) = [eIF4F]_0, [AUG](t) = \frac{k_2 \cdot [eIF4F]_0}{k_3([60S]_0 - [80S](t))}. \end{aligned} \quad (37)$$

From these equations, one can determine the effective duration of the Stage 1: by definition, it will be finished when one of the two conditions ($[40S] \gg [eIF4F]$, $[60S] \gg [AUG]$) will be violated, which happens at times $\sim \frac{[40S]_0}{k_2 \cdot [eIF4F]_0}$ and $\sim \frac{[60S]_0}{k_2 \cdot [eIF4F]_0}$ correspondingly, hence, the second condition will be violated first (from $[60S]_0 < [40S]_0$).

3.3.4 Stage 2: first stage of protein elongation, initiation is still rapid

The Stage 2 is characterized by conditions $[eIF4F] \ll [40S]$, $[60S] \ll [AUG]$. This fact can be used for deriving the quasi-steady state approximation: we assume that the reaction fluxes in two network cycles (R1-R2 and R3-R4) are independently balanced:

$$k_1[40S] \cdot [eIF4F] = k_2[mRNA : 40S], k_3[AUG] \cdot [60S] = k_4[80S]. \quad (38)$$

Then (22) is simplified and, using the conservation laws, we have a single equation on $[40S]$:

$$\frac{d[40S](t)}{dt} = \frac{-k_2[eIF4F]_0 \cdot [40S](t)}{\frac{k_2}{k_1} + [40S](t)} + \frac{k_4[60S]_0(A - [40S](t))}{\frac{k_4}{k_3} + A - [40S](t)}, \quad (39)$$

where $A = [40S](t) + [AUG](t)$ is a constant quantity conserved accordingly to the quasi-steady state approximation. Equation (39) can be already integrated but let us further simplify it for our analysis. Having in mind $k_4 \ll k_3$ and assuming that at the beginning of the Stage 2 $[AUG] \gg \frac{k_4}{k_3}$, we can simplify (39) to

$$\frac{d[40S](t)}{dt} = \frac{-k_2[eIF4F]_0 \cdot [40S](t)}{\frac{k_2}{k_1} + [40S](t)} + k_4[60S]_0, \quad (40)$$

and, further, assuming that at the beginning of the Stage 2 we have $[40S] \gg [AUG]$ let us approximate the right-hand side of the equation by a piecewise-linear function

$$\frac{d[40S](t)}{dt} \approx \begin{cases} -k_2[eIF4F]_0 \frac{\frac{k_2}{k_1} + [40S]|_{t=t''}}{3\frac{k_2}{k_1} + [40S]|_{t=t''}} + k_4[60S]_0, & \text{if } [40S](t) > \frac{k_2}{k_1}, \\ -\frac{k_1[eIF4F]_0}{2} \cdot [40S](t) + k_4[60S]_0, & \text{if } [40S](t) < \frac{k_2}{k_1}, \end{cases} \quad (41)$$

where $[40S]|_{t=t''}$ is the amount of $40S$ at the beginning of the Stage 2. Then the descent of $[40S](t)$ can be separated into linear and exponential phases:

$$[40S](t) \approx \begin{cases} [40S]|_{t=t''} - K_1 \cdot (t - t''), & \text{if } t < t'' + \tau_2 \\ [40S]_{s2} - ([40S]_{s2} - [40S]|_{t=t''}) \exp(-K_2(t - t'' - \tau_2)), & \text{if } t > t'' + \tau_2 \end{cases}, \quad (42)$$

where K_1, K_2 are linear and exponential slopes and $[40S]_{s2}$ is the quasi-steady state value of $[40S]$ at the end of the Stage 2:

$$\begin{aligned}
K_1 &= k_2 eIF4F_0 \frac{\frac{k_2}{k_1} + [40S]_{t=t''}}{3 \frac{k_2}{k_1} + [40S]_{t=t''}} - k_4 [60S]_0 \\
K_2 &= \frac{k_1 [eIF4F]_0}{2} \\
[40S]_{s2} &= \frac{k_2}{k_1} \frac{1}{\frac{k_2 [eIF4F]_0}{k_4 [60S]_0} - 1} \\
\tau_2 &= \frac{[40S]_{t=t_2} - \frac{k_2}{k_1}}{K_1}
\end{aligned} \tag{43}$$

Other dynamic variables are expressed through $[40S](t)$ as

$$\begin{aligned}
[eIF4F](t) &= [eIF4F]_0 \frac{\frac{k_2}{k_1}}{\frac{k_2}{k_1} + [40S](t)} \\
[mRNA : 40S](t) &= [eIF4F]_0 \cdot \frac{[40S](t)}{\frac{k_2}{k_1} + [40S](t)} \\
[AUG](t) &= [40S]_{t=t''} + \frac{k_2 [eIF4F]_0}{k_3 ([60S]_0 - [40S]_0 + [40S]_{t=t''})} - [40S](t). \\
[60S](t) &= [60S]_0 \frac{\frac{k_4}{k_3}}{\frac{k_4}{k_3} + [AUG](t)} \\
[80S]_0(t) &= [60S]_0 \frac{[AUG](t)}{\frac{k_4}{k_3} + [AUG](t)}
\end{aligned} \tag{44}$$

At some point, the amount of free small ribosomal subunit 40S, which is abundant at the beginning of the Stage 2, will not be sufficient to support rapid translation initiation. Then the initiation factor $eIF4F$ will not be the limiting factor in the initiation and the condition $[40S] \gg [eIF4F]$ will be violated. We can estimate this time as $t''' = \frac{[40S]_0}{k_2 \cdot [eIF4F]_0}$.

3.3.5 Stage 3: steady protein elongation, speed of initiation equals to speed of elongation

During the Stage 3 all fluxes in the network become balanced and the translation arrives at the steady state. From Fig. 5 it is clear that the relaxation goes independently in the cycle $R3 - R4$, where the relaxation equations are simply

$$[60S](t) = [60S]_s + ([60S]_{t=t'''} - [60S]_s) e^{-(k_4 + k_3 [AUG]_{t=t'''})(t-t''')}, [80S](t) = [60S]_0 - [60S](t). \tag{45}$$

where t''' is the time when the Stage 3 of the relaxation starts. This relaxation goes relatively fast, since $k_3 [AUG]_{t=t'''}$ is relatively big. So, during the Stage 3, one can consider the cycle $R1 - R4$ equilibrated, with $[80S] = [80]_s$, $[60S] = [60]_s$ values.

Hence, the relaxation during the Stage 3 consists in redistributing concentrations of 40S and mRNA:40S to their steady states in a linear chain of reactions $R1 - R2$ (the value of $[AUG]$ is relatively big and can be adjusted from the conservation law (23)). Using the pseudo-linear approximation of this stage (see Fig.5), we can easily write down the corresponding approximate relaxation equations:

$$\begin{aligned}
[mRNA : 40S](t) &= \\
[mRNA : 40]_s (1 - e^{-k_2 t}) + B (e^{-k_1 [eIF4F]_{t=t'''}(t-t''')} - e^{-k_2 (t-t''')}) + [mRNA : 40S]_{t=t'''} e^{-k_2 (t-t''')}, \\
[40S](t) &= [40S]_s + B e^{-k_2 (t-t''')}
\end{aligned} \tag{46}$$

where $B = (1 - \frac{k_2}{k_1[eIF4F]_{t=t'''}})([40]_s - [40S]_{t=t'''})$. $[40]_s$ and $[mRNA : 40S]_s$ are the steady-state values of the corresponding variables, see (28). The values $[60S]_{t=t'''}$, $[eIF4F]_{t=t'''}$, $[AUG]_{t=t'''}$ and $[mRNA : 40S]_{t=t'''}$ can be estimated from (44), using the $[40S]_{t=t'''}$ value. The relaxation time at this stage equals

$$\tau_3 = \max\left(\frac{1}{k_1[eIF4F]_{t=t'''}} , \frac{1}{k_2} , \frac{1}{k_4 + k_3[AUG]_{t=t'''}}\right)$$

The solution for the Stage 3 can be further simplified if $k_2 \ll k_1[eIF4F]_{t=t'''}$ or $k_2 \gg k_1[eIF4F]_{t=t'''}$.

3.3.6 Transitions between stages

Along the trajectory of the dynamical system (22) there are three dominant system each one transforming into an other. At the transition between stages, two neighbor dominant systems are united and then split. Theoretically, there might be situations when the system can stay in these transition zones for long periods of time, even infinitely. However, in the model (22) this is not the case: the trajectory rapidly passes through the transition stages and jumps into the next dominant system approximation.

Three dominant approximations can be glued, using the concentration values at the times of the switching of dominant approximation as initial values for the next stage. Note that the Stage 2 has essentially one degree of freedom since it can be approximated by a single equation (39). Hence, one should only know one initial value $[40S]_{t=t''}$ to glue the Stages 1 and 2. The same is applied to the gluing of Stages 2 and 3, since in the end of Stage 2 all variable values are determined by the value of $[40S]_{t=t''}$.

3.3.7 Case of always limiting initiation

As it follows from our analysis, the most critical parameter of the non-linear protein translation model is the ratio $\beta = \frac{k_2[eIF4F]_0}{k_4[60S]_0}$. Above we have considered the case $\beta > 1$ which is characterized by a switch of the limiting factor in the initiation (from $eIF4F$ at the Stages 1 and 2 to $40S$ at the Stage 3).

In the case $\beta < 1$ the dynamics becomes simpler and consists of one single stage: relaxation accordingly to (36) and further with correction (37) with the relaxation time $\sim \frac{1}{k_4}$ (the quasiequilibrium approximation corresponding to the Stage 1 works well for the whole translation process). The reason for this is that if the initiation is not efficient then the system is never in the situation of the Stage 2 conditions when the cycle R1-R2 is balanced with much bigger flux than the cycle R3-R4. This approximation is the more exact the smaller β value, however, the value of β should not be necessary very small. For example, for the default parameter values of the model $\beta = 0.48$, and it well reproduces the dynamics (see Fig. 7c-d). From numerical experiments one can see that even for $\beta = 0.95$ the dynamics is qualitatively well reproduced.

To model the A-cap structure effect with very weak capacity for initiation (assembly of the initiation factors and $40S$ subunit), we should also consider the case

$$k_1 \ll k_4 \ll k_2 \ll k_3, \quad (47)$$

for which the solution derived above is not directly applicable. However, the analytical calculations in this case can be performed in the same fashion as above. The detailed derivation of the solution is given in the Supplementary materials. The effect of putting k_1 very small on the steady state protein synthesis and the relaxation time is shown on Fig. 7.

In a similar way all possible solutions of the equations (22) with very strong inhibitory effect of microRNA on a particular translation step can be derived. These solutions will describe the situation when the effect of microRNA is so strong that it changes the dominant system (limiting place of the network) by violating the initial constraints (26) on the parameters (for example, by making k_3 smaller than other k_i s). Such possibility exists, however, it can require too strong (non-physiological) effect of microRNA-dependent translation inhibition.

3.3.8 Effect of microRNA on the translation dynamics

As a result of the above analysis, we can assemble an approximate solution of the non-linear system under assumptions (26) about the parameters. An example of the approximate solution is given on




Figure 7: Examples of the exact (circles) and approximate (solid lines) solutions of the non-linear protein translation model. a) For our set of parameters $k_1 = 2$, $k_2 = 3$, $k_3 = 50$, $k_4 = 0.1$; b) For parameters $k_1 = 1$, $k_2 = 5$, $k_3 = 50$, $k_4 = 0.01$; c) For the set of parameters from [37]; d) Reaction fluxes for the set of parameters c). Dashed black vertical lines denote evaluated transition points between the dynamics stages. Dashed red vertical points denote the time points where $[40S](t) = 10 \cdot [eIF4F](t)$ and $[40S](t) = [eIF4F](t)/10$ respectively.

Fig. 7. The advantage of such a semi-analytical solution is that one can predict the effect of changing the system parameters. For example, on Fig. 7b the solution is compared to an exact numerical one, where the parameters have been changed but still obey the initial constraints (26).

One of the obvious predictions is that the dynamics of the system is not sensitive to variations of k_3 , so if microRNA acts on the translation stage controlled by k_3 then no microRNA effect could be observed looking at the system dynamics (being the fastest one, k_3 is not a critical parameter in any scenario).

If microRNA acts on the translation stage controlled by k_4 (for example, by ribosome stalling mechanism) then we should consider two cases of efficient ($\beta > 1$) and inefficient ($\beta < 1$) initiation. In the first case the steady state protein synthesis rate is controlled by k_4 (as the slowest, limiting step) and any effect on k_4 would lead to the proportional change in the steady state of protein production. By contrast, in the case of inefficient initiation, the steady state protein synthesis is not affected by k_4 . Instead, the relaxation time is affected, being $\sim \frac{1}{k_4}$. However, diminishing k_4 increases the β parameter, hence, this changes “inefficient initiation” scenario for the opposite, hence, making k_4 critical for the steady state protein synthesis anyway when k_4 becomes smaller than $\frac{k_2[eIF4F]_0}{[60S]_0}$. For example, for the default parameters of the model, decreasing k_4 value first leads to no change in the steady state rate of protein synthesis but increases the relaxation time and, second, after the threshold value $\frac{k_2[eIF4F]_0}{[60S]_0}$ starts to affect the steady state protein synthesis rate directly (see Fig. 8A). This is in contradiction to the message from [37] that the change in k_4 by several orders of magnitude does not change the steady state rate of protein synthesis.

Analogously, decreasing the value of k_2 can convert the “efficient” initiation scenario into the opposite after the threshold value $\frac{k_4[60S]_0}{[eIF4F]_0}$. We can recapitulate the effect of decreasing k_2 in the following way. 1) in the case of efficient initiation k_2 does not affect the steady state protein synthesis rate up to the threshold value after which it affects it in a proportional way. The relaxation time drastically increases, because decreasing k_2 leads to elongation of all dynamical stages durations (for example, we have estimated the time of the end of the dynamical Stage 2 as $t''' = \frac{[40S]_0}{k_2[eIF4F]_0}$). However, after the threshold value the relaxation time decreases together with k_2 , quickly dropping to its unperturbed value. 2) in the case of “inefficient” initiation the steady state protein synthesis rate depends proportionally on the value of k_2 (31), while the relaxation time is not affected (see Fig. 8).

MicroRNA action on k_1 directly does not produce any strong effect neither on the relaxation time nor on the steady state protein synthesis rate. This is why in the original work [37] cap-dependent mechanism of microRNA action was taken into account through effective change of the $[eIF4F]_0$ value (total concentration of the translation initiation factors), which is a critical parameter of the model (see 31).

The effect of microRNA on various mechanism and in various experimental settings (excess or deficit of eIF4F, normal cap or A-cap) is recapitulated in Table 2. The conclusion that can be made from this table is that all four mechanism shows clearly different pattern of behaviour in various experimental settings. From the simulations one can make a conclusion that it is still not possible to distinguish between the situation when microRNA does not have any effect on protein translation and the situation when it acts on the step which is neither rate limiting nor ‘second rate limiting’ in any experimental setting (k_3 in our case). Nevertheless, if any change in the steady-state protein synthesis or the relaxation time is observed, theoretically, it will be possible to specify the mechanism responsible for it.

4 Discussion

The role of microRNA in gene expression regulation is discovered and confirmed since ten years, however, there is still a lot of controversial results regarding the role of concrete mechanisms of microRNA-mediated protein synthesis repression. Some authors argue that it is possible that the different modes of microRNA action reflect different interpretations and experimental approaches, but the possibility that microRNAs do indeed silence gene expression via multiple mechanisms also exists. Finally, microRNAs might silence gene expression by a common and unique mechanism; and the multiple modes of action represent secondary effects of this primary event [6, 9, 11].

The main reason for accepting a possible experimental bias could be the studies in vitro, where




Figure 8: Effect of decreasing some model parameters mimicking different mechanisms of miRNA action on translation. Relaxation time here is defined as the latest time at which any chemical species in the model differs from its final steady state by 10%. A) and B) correspond to the scenario with "inefficient" initiation, with use of the model parameters proposed in [37] ($k_1 = k_2 = 2$, $k_3 = 5$, $k_4 = 1$, $[eIF4F]_0 = 6$, $[60S]_0 = 25$, $[40S]_0 = 100$), which gives $\beta = 0.48 < 1$. C) and D) correspond to the scenario with "efficient" initiation, with our choice of parameters ($k_1 = 2$, $k_2 = 3$, $k_3 = 50$, $k_4 = 0.1$, $[eIF4F]_0 = 6$, $[60S]_0 = 25$, $[40S]_0 = 100$), which gives $\beta = 7.2 > 1$. The absciss value indicates the degree of inhibition (decreasing) of a parameter. E-H) same as A-D) but for a modified cap structure, modeled by reduced k_1 parameter: $k_1 = 0.01$ for these curves, the other parameters are the same as on A-D) correspondingly.

Table 2: Modeling of four mechanisms of microRNA action in the non-linear protein translation model. MicroRNA action effect is described for the protein synthesis steady rate and the relaxation time (see also Fig. 8). It is assumed that the ribosome assembly step in protein translation, described by the k_3 rate constant, is not rate limiting

Observable value	Initiation(k_1)	Step after initiation(k_2)	Ribosome assembly (k_3)	Elongation (k_4)
Wild-type cap, inefficient initiation				
<i>Steady-state rate</i>	slightly decreases	decreases	no change	decreases after threshold
<i>Relaxation time</i>	no change	no change	no change	goes up and down
Wild-type cap, efficient initiation				
<i>Steady-state rate</i>	no change	slightly decreases after strong inhibition	no change	decreases
<i>Relaxation time</i>	no change	goes up and down	no change	no change
A-cap, inefficient initiation				
<i>Steady-state rate</i>	decreases	decreases	no change	slightly decreases after strong inhibition
<i>Relaxation time</i>	no change	no change	no change	goes up and down
A-cap, efficient initiation				
<i>Steady-state rate</i>	decreases after threshold	slightly decreases after strong inhibition	no change	decreases
<i>Relaxation time</i>	goes up and down	goes up and down	no change	increases

conditions are strongly different from situation in vivo. Indeed, inside the cell, mRNAs (microRNA targets) exist as ribonucleoprotein particles or mRNPs, and second, all proteins normally associated with mRNAs transcribed in vivo are absent or at least much different from that bound to the same mRNA in an in vitro system or following the microRNAs transfection into cultured cells. The fact that RNA-binding proteins strongly influence the final outcome of microRNA regulation is proved now by several studies [3, 12, 22].

The mathematical results provided in this paper suggests a complementary view on the co-existence of multiple microRNA-mediated mechanisms of translation repression. Mathematical modeling suggests to us to ask a question: *if multiple mechanisms act at the same time, would all of them equally contribute to the final observable repression of protein synthesis or its dynamics?* The dynamical limitation theory gives an answer: *the effect of microRNA action will be observable and measurable in two cases: 1) if it affects the dominant system of the protein translationary machinery, or 2) if the effect of microRNA action is so strong that it changes the limiting place (the dominant system).*

In a limited sense, this means, in particular, that the protein synthesis steady rate is determined by the limiting step in the translation process and any effect of microRNA will be measurable only if it affects the limiting step in translation, as it was demonstrated in [37]. Due to the variety of external conditions, cellular contexts and experimental systems the limiting step in principle can be any in the sequence of events in protein translation, hence, this or that microRNA mechanism can become dominant in a concrete environment. However, when put on the language of equations, the previous statement already becomes non-trivial in the case of non-linear dynamical models of translation (and even linear reaction networks with non-trivial network structure). Our analysis demonstrates that the limiting step in translation can change with time, depends on the initial conditions and is not represented by a single reaction rate constant but rather by some combination of several model parameters. Methodology of dynamical limitation theory that we had developed [16, 17], allows to deal with these situations on a solid theoretical ground.

Furthermore, in the dynamical limitation theory, we generalize the notion of the limiting step to the notion of dominant system, and this gives us a possibility to consider not only the steady state rate but also some dynamical features of the system under study. One of the simplest measurable dynamical feature is the *protein synthesis relaxation time*, i.e. the time needed for protein synthesis to achieve its steady state rate. The general idea of “relaxation spectrometry” goes back to the works of Manfred Eigen, a Nobel laureate [8] and is still underestimated in systems biology. Calculation of the relaxation time (or times) requires careful analysis of time scales in the dynamical system, which is greatly facilitated by the recipes proposed in [17, 45]. As we have demonstrated in our semi-analytical solutions, measuring the steady state rate and relaxation time at the same time allows to detect which step is possibly affected by the action of microRNA (resulting in effective slowing down of this step). To our knowledge, this idea was never considered before in the studies of microRNA-dependent expression regulation. The table 2 recapitulates predictions allowing to discriminate a particular mechanism of microRNA action.

Thus, analysis of the transient dynamics gives enough information to verify or reject a hypothesis about a particular molecular mechanism of microRNA action on protein translation. For multiscale systems only that action of microRNA is distinguishable which affects the parameters of dominant system (critical parameters), or changes the dominant system itself. Dominant systems generalize and further develop the old and very popular idea of limiting step. Algorithms for identifying dominant systems in multiscale kinetic models are straightforward but not trivial and depend only on the ordering of the model parameters but not on their concrete values. Asymptotic approach to kinetic models of biological networks suggest new directions of thinking on a biological problem, making the mathematical model a useful tool accompanying biological reasoning and allowing to put in order diverse experimental observations.

5 Author contributions

AZ and AG have written the main body of the manuscript. NM, NN and AHB provided the critical review of miRNA mechanisms and contributed to writing the manuscript. AG and AZ developed the mathematical methodology for identifying dominant systems. AZ performed the analytical computations and numerical simulations. All authors have participated in discussing the results and model predictions.

6 Acknowledgements

We acknowledge support from Agence Nationale de la Recherche (ANR CALAMAR project) and from the Projet Incitatif Collaboratif “Bioinformatics and Biostatistics of Cancer” at Institut Curie. AZ and EB are members of the team “Systems Biology of Cancer” Equipe labellisée par la Ligue Nationale Contre le Cancer. We thank Vitaly Volpert and Laurence Calzone for inspiring and useful discussions.

References

- [1] Baillat, D. and R. Shiekhataar, Functional dissection of the human TNRC6 (GW182-related) family of proteins. *Mol Cell Biol*, 2009. 29(15): p. 4144-55.
- [2] Bandres, E., et al., Identification by Real-time PCR of 13 mature microRNAs differentially expressed in colorectal cancer and non-tumoral tissues. *Mol Cancer*, 2006. 5: p. 29.
- [3] Bhattacharyya, S.N., et al., Relief of microRNA-mediated translational repression in human cells subjected to stress. *Cell*, 2006. 125(6): p. 1111-24.
- [4] Bykov, V., Goldfarb, I., Gol'dshtein, V., Maas, U. (2006). On a Modified Version of ILDM Approach: Asymptotical Analysis Based on Integral Manifolds Method, *IMA J. of Applied Mathematics* 71 (3), 359–382.
- [5] Boyd, R. K. (1978). Some common oversimplifications in teaching chemical kinetics, *J. Chem. Educ.* 55, 84–89.
- [6] Chekulaeva, M., and Filipowicz, W. (2009). Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells. *Curr Opin Cell Biol* 21, 452-460.
- [7] Chendrimada, T.P., et al., MicroRNA silencing through RISC recruitment of eIF6. *Nature*, 2007. 447(7146): p. 823-8.
- [8] Eigen, M. (1972). Immeasurably fast reactions, Nobel Lecture, December 11, 1967, In: *Nobel Lectures, Chemistry* 1963-1970, Amsterdam: Elsevier, 170203.
- [9] Eulalio A., Huntzinger E. and Izaurralde E. (2008) Getting to the root of miRNA-mediated gene silencing. *Cell* **132**(1):p.9-14.
- [10] Eulalio, A., E. Huntzinger, and E. Izaurralde. (2008) GW182 interaction with Argonaute is essential for miRNA-mediated translational repression and mRNA decay. *Nat Struct Mol Biol* 15(4): p. 346-53.
- [11] Filipowicz, W., S.N. Bhattacharyya, and N. Sonenberg. (2008) Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet.* **9**(2): p. 102-14.
- [12] George, A.D. and Tenenbaum, S.A. (2006) MicroRNA modulation of RNA-binding protein regulatory elements. *RNA Biol*, 3, 57-59.
- [13] Gorban, A.N., Karlin, I.V., Zinovyev, A.Yu. (2004). Invariant grids for reaction kinetics, *Physica A* 333, 106–154. Preprint online: <http://arxiv.org/abs/cond-mat/0307076>.
- [14] Gorban, A. N., & Karlin, I. V. (2005). *Invariant manifolds for physical and chemical kinetics*, volume 660 of *Lect. Notes Phys.* Berlin–Heidelberg–New York: Springer.
- [15] Gorban, A. N., & Radulescu, O. (2007). Dynamical robustness of biological networks with hierarchical distribution of time scales, *IET Syst. Biol.*, 1, 238-246.
- [16] Gorban, A. N. & Radulescu, O. (2008). Dynamic and static limitation in reaction networks, revisited, *Advances in Chemical Engineering*, 34, 103-173; e-print: <http://arxiv.org/abs/physics/0703278>
- [17] Gorban A., Radulescu O., Zinovyev A. (2009). Asymptotology of chemical reaction networks. *Chemical Engineering Science*, in press, doi:10.1016/j.ces.2009.09.005. Arxive: <http://arxiv.org/abs/0903.5072>.
- [18] Gu, S., et al., Biological basis for restriction of microRNA targets to the 3' untranslated region in mammalian mRNAs. *Nat Struct Mol Biol*, 2009. 16(2): p. 144-50.

- [19] Huang, J., et al., Derepression of microRNA-mediated protein translation inhibition by apolipoprotein B mRNA-editing enzyme catalytic polypeptide-like 3G (APOBEC3G) and its family members. *J Biol Chem*, 2007. 282(46): p. 33632-40.
- [20] Humphreys, D.T., et al., MicroRNAs control translation initiation by inhibiting eukaryotic initiation factor 4E/cap and poly(A) tail function. *Proc Natl Acad Sci U S A*, 2005. 102(47): p. 16961-6.
- [21] Jackson, R.J., and Standart, N. (2007). How do microRNAs regulate gene expression? *Sci STKE*. 2007 Jan 2;2007(367):re1.
- [22] Jing, Q., Huang, S., Guth, S., Zarubin, T., Motoyama, A., Chen, J., Di Padova, F., Lin, S.C., Gram, H. and Han, J. (2005) Involvement of microRNA in AU-rich element-mediated mRNA instability. *Cell* 120, 623-634.
- [23] Johnston, H. S. (1966). *Gas phase reaction theory*, New York: Roland Press.
- [24] Kazantzis, N. & Kravaris, C. (2006). A New Model Reduction Method for Nonlinear Dynamical Systems using Singular PDE Theory, In: *Model Reduction and Coarse-Graining Approaches for Multiscale Phenomena*, A.N. Gorban, N. Kazantzis, Y.G. Kevrekidis, H.C. Ottinger and C. Theodoropoulos (eds.), Springer, 3–15.
- [25] Karaa, Z.S., et al., The VEGF IREs are differentially susceptible to translation inhibition by miR-16. *RNA*, 2009. 15(2): p. 249-54.
- [26] Kim, D.H., et al., MicroRNA-directed transcriptional gene silencing in mammalian cells. *Proc Natl Acad Sci U S A*, 2008. 105(42): p. 16230-5.
- [27] Kiriakidou, M., et al., An mRNA m7G cap binding-like motif within human Ago2 represses translation. *Cell*, 2007. 129(6): p. 1141-51.
- [28] Kong, Y.W., et al. (2008) The mechanism of micro-RNA-mediated translation repression is determined by the promoter of the target gene. *Proc Natl Acad Sci U S A* 105(26): p. 8866-71.
- [29] Kozak, M. (2008). Faulty old ideas about translational regulation paved the way for current confusion about how microRNAs function. *Gene* 423, 108-115.
- [30] Kruskal M.D. Asymptotology. *Proceedings of Conference on Mathematical Models on Physical Sciences*. Englewood Cliffs, HJ: Prentice-Hall, 1963, 1748.
- [31] Lam, S. H. (1993). Using CSP to Understand Complex Chemical Kinetics, *Combustion Science and Technology*, 89 (5), 375-404.
- [32] Lam, S. H., & Goussis, D. A. (1994). The CSP Method for Simplifying Kinetics, *International Journal of Chemical Kinetics* 26, 461–486.
- [33] Lytle, J.R., T.A. Yario, and J.A. Steitz, Target mRNAs are repressed as efficiently by microRNA-binding sites in the 5' UTR as in the 3' UTR. *Proc Natl Acad Sci U S A*, 2007. 104(23): p. 9667-72.
- [34] Maas, U., & Pope, S.B. (1992). Simplifying chemical kinetics: intrinsic low – dimensional manifolds in composition space, *Combustion and Flame* 88, 239–264.
- [35] Maroney, P.A., et al., Evidence that microRNAs are associated with translating messenger RNAs in human cells. *Nat Struct Mol Biol*, 2006. 13(12): p. 1102-7.
- [36] Mathonnet G., Fabian M.R., Svitkin Y.V. et al. (2007). MicroRNA inhibition of translation initiation in vitro by targeting the cap-binding complex eIF4F. *Science* **317**: 1764-1767.
- [37] Nissan T. and Parker R. (2008). Computational analysis of miRNA-mediated repression of translation: Implications for models of translation initiation inhibition. *RNA* **14**, 1480-1491.
- [38] Nottrott, S., M.J. Simard, and J.D. Richter, Human let-7a miRNA blocks protein production on actively translating polyribosomes. *Nat Struct Mol Biol*, 2006. 13(12): p. 1108-14.
- [39] Olsen, P.H. and V. Ambros, The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the initiation of translation. *Dev Biol*, 1999. 216(2): p. 671-80.
- [40] Orom, U.A., F.C. Nielsen, and A.H. Lund, MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation. *Mol Cell*, 2008. 30(4): p. 460-71.

- [41] Place, R.F., et al., MicroRNA-373 induces expression of genes with complementary promoter sequences. *Proc Natl Acad Sci U S A*, 2008. 105(5): p. 1608-13.
- [42] Pillai, R.S., et al., Inhibition of translational initiation by Let-7 MicroRNA in human cells. *Science*, 2005. 309(5740): p. 1573-6.
- [43] Petersen, C.P., et al., Short RNAs repress translation after initiation in mammalian cells. *Mol Cell*, 2006. 21(4): p. 533-42.
- [44] Rate-controlling step (2007). In: IUPAC Compendium of Chemical Terminology, E-version, <http://goldbook.iupac.org/R05139.html>.
- [45] Radulescu O., Gorban A., Zinovyev A., Lilienbaum, A. (2008). Robust simplifications of multiscale biochemical networks, *BMC Systems Biology*, **2** (1):86
- [46] Roussel, M.R., & Fraser, S.J. (1991). On the geometry of transient relaxation. *J. Chem. Phys.* **94**, 7106–7111.
- [47] Temkin, O.N., Zeigarnik, A.V., & Bonchev, D.G. (1996). *Chemical Reaction Networks: A Graph-Theoretical Approach*, Boca Raton, FL: CRC Press.
- [48] Thermann R. and Hentze M.W. (2007). Drosophila miR2 induces pseudo-polysomes and inhibits translation initiation. *Nature* **447**: 875-878.
- [49] Valencia-Sanchez, M.A., Liu, J., Hannon, G.J., and Parker, R. (2006). Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev* **20**, 515-524.
- [50] Vasudevan, S., Y. Tong, and J.A. Steitz, Switching from repression to activation: microRNAs can up-regulate translation. *Science*, 2007. 318(5858): p. 1931-4.
- [51] Wakiyama, M., et al., Let-7 microRNA-mediated mRNA deadenylation and translational repression in a mammalian cell-free system. *Genes Dev*, 2007. 21(15): p. 1857-62.
- [52] Wang, B., A. Yanez, and C.D. Novina, MicroRNA-repressed mRNAs contain 40S but not 60S components. *Proc Natl Acad Sci U S A*, 2008. 105(14): p. 5343-8.
- [53] Wang, B., et al., Recapitulation of short RNA-directed translational gene silencing in vitro. *Mol Cell*, 2006. 22(4): p. 553-60.
- [54] White, R. B. (2006). *Asymptotic Analysis of Differential Equations*, London: Imperial College Press & World Scientific.
- [55] Yablonsky, G.S., Olea, M., & Marin, G.B. (2003). Temporal Analysis of Products (TAP): Basic Principles, Applications and Theory, *Journal of Catalysis* **216**, 120–134.
- [56] Yekta S., Shih I.H., Bartel D.P. (2004) MicroRNA-directed cleavage of HOXB8 mRNA. *Science* **304**(5670):594-596.
- [57] Zagaris, A., Kaper, H.G., Kaper, T.J. (2004). Analysis of the computational singular perturbation reduction method for chemical kinetics, *J. Nonlinear Sci.* **14**, 5991.